

Rekommenderad komposition av vätsketerapi under cisplatininfusion. VSTB:s rekommendation 2015-09-16

	<u>Vätska</u>	<u>Tillsatser</u>	<u>Tid</u>
Prehydrering:	Glucos 25mg/mL buffrad (Na = 70 mmol/L) 125 mL/m ² /tim ≥750 mL/m ²	K ⁺ 20 mmol/L Mg ⁺⁺ 5 mmol/L Ca ⁺⁺ 2,3 mmol/L ¹	6 - 12 tim
(Cisplatin:	100 - 1000 ml NaCl		1 – 48 tim ²)
Under cisplatin: (parallell infusion)	Glucos 25mg/mL buffrad (Na = 70 mmol/L) 125 mL/m ² /tim 3000 mL/m ² /24 tim	K ⁺ 20 mmol/L Mg ⁺⁺ 5 mmol/L Ca ⁺⁺ 2,3 mmol/L ¹ Mannitol 150 mg/mL, 100 mL/L = 15 g/L	1 – 48 tim
Efter cisplatin:	Glucos 25mg/mL buffrad (Na = 70 mmol/L) 125 mL/m ² /tim 3000 mL/m ² /24 tim	K ⁺ 20 mmol/L Mg ⁺⁺ 5 mmol/L Ca ⁺⁺ 2,3 mmol/L ¹	24 tim

¹ 10 mL Calcium gluconate 10%, B. Braun (licenspreparat), eller motsv./1000 mL infusionslösning

² Protokollberoende infusionstid